

## Unit - 6 - Numerical Integration

### \* Task - 1: Trapezoidal Rule

1 State Trapezoidal rule with  $n=10$  and evaluate  $\int_0^1 e^x \cdot dx$ .

=> For  $n=10$ , Trapezoidal Rule

$$\int_a^b f(x) \cdot dx = \frac{h}{2} [ (y_0 + y_n) + 2(y_1 + y_2 + y_3 + y_4 + y_5 + y_6 + y_7 + y_8) ]$$

=> Let (Given  $f(x) = e^x$  and  $b=1$  and  $a=0$ )

So,  $h = \frac{b-a}{n} = \frac{1-0}{10} = 0.1$

X	0	0.1	0.2	0.3	0.4	0.5
Y	1	1.1052	1.2214	1.3499	1.4912	1.6487

X	0.6	0.7	0.8	0.9	0.1
Y	1.8221	2.0137	2.2255	2.4596	2.7183

By Trapezoidal Rule,

$$\int_0^1 e^x \cdot dx = \frac{h}{2} [Y_0 + Y_{10} + 2(Y_1 + Y_2 + Y_3 + Y_4 + Y_5 + Y_6 + Y_7 + Y_8 + Y_9)]$$

$$= \frac{0.1}{2} [(1 + 2.7183) + 2(1.1052 + 1.2214 + 1.3499 + 1.4918 + 1.6487 + 1.8221 + 2.0137 + 2.2255 + 2.4596)]$$

$$= 0.05 [3.7183 + 2(15.3379)]$$

$$\int_0^1 e^x \cdot dx = 1.7197$$

2 Evaluate  $\int_0^1 e^{-x^2} \cdot dx$  Using trapezoidal rule with  $h=0.1$

Here, Given  $F(x) = e^{-x^2}$ ,  $a=0$ ,  $b=1$ ,  $h=0.1$

$$\text{So, } n = \frac{b-a}{h} = \frac{1-0}{0.1} = 10$$

X	0	0.1	0.2	0.3	0.4	0.5
Y	1	0.99	0.9608	0.9139	0.8521	0.7788

X	0.6	0.7	0.8	0.9	1
Y	0.6977	0.6126	0.5273	0.4449	0.3679

By Trapezoidal Rule,

$$\int_0^1 e^{-x^2} dx = \frac{0.1}{2} [ (1 + 0.3679) + 2(0.99 + 0.9608 + 0.9139 + 0.8521 + 0.7788 + 0.6977 + 0.6126 + 0.5273 + 0.4449) ]$$

$$= 0.05 [ 1.3679 + 2(6.7781) ]$$

$$\int_0^1 e^{-x^2} dx = 0.7462$$

3 Evaluate  $\int_0^1 \frac{1}{1+x^2} dx$  Using

trapezoidal rule with  $h = 0.2$

$\Rightarrow$  Here, Given  $f(x) = \frac{1}{1+x^2}$  and

$a = 0$ ,  $b = 1$  and  $h = 0.2$

$$\text{So, } n = \frac{b-a}{h} = \frac{1}{0.2} = 5$$

x	0	0.2	0.4	0.6	0.8	<del>1.0</del> 1
y	1	0.9615	0.8621	0.7353	0.6097	0.5

By Trapezoidal Rule,

$$\int_0^1 \frac{1}{1+x^2} \cdot dx = 0.2 \left[ (1 + 0.5) + 2(0.9615 + 0.8621 + 0.7353 + 0.6097) \right]$$

$$\int_0^1 \frac{1}{1+x^2} \cdot dx = 0.78372$$

4 Use trapezoidal rule to evaluate

$\int_0^2 \frac{x}{\sqrt{2+x^2}} \cdot dx$  dividing the interval into four equal parts.

$$\Rightarrow \text{Let, Given } f(x) = \int_0^2 \frac{x}{\sqrt{2+x^2}},$$

$$a = 0, \quad b = 2 \quad \text{and} \quad n = 4,$$

$$\text{So, } h = \frac{b-a}{n} = \frac{2}{4} = 0.5$$

X	0	0.5	1	1.5	2
Y	0	0.3333	0.5774	0.7276	0.8165

By Trapezoidal Rule,

$$\int_0^2 \frac{x}{\sqrt{2+x^2}} dx = \frac{h}{2} [y_0 + y_4 + 2(y_1 + y_2 + y_3)]$$

$$= \frac{0.5}{2} [0 + 0.8165 + 2(0.3333 + 0.5774 + 0.7276)]$$

$$= 1.0233$$

5 Find the area bounded by the curve  $Y = f(x)$  and the X-axis from  $x = 7.47$  to  $x = 7.52$  from the following Table.

X	7.47	7.48	7.49	7.50	7.51	7.52
f(x)	1.93	1.95	1.98	2.01	2.03	2.06

$\Rightarrow$  Here, Given  $n = 5$ ,  $a = 7.47$  and  $b = 7.52$

So,  $h = \frac{b-a}{n} = \frac{7.52-7.47}{5} = 0.01$

By Trapezoidal Rule,

$$\int_a^b f(x) \cdot dx = \frac{0.01}{2} [1.93 + 2.06 + 2(1.95 + 1.98 + 2.01 + 2.03)]$$

$$\int_a^b f(x) \cdot dx = 0.09965$$

Brain Spot

\* Task: 2 : Simpson's  $\frac{1}{3}$  rule, Simpson's  $\frac{3}{8}$  rule and Weddle's rule.

1 Evaluate  $\int_0^6 \frac{1}{1+x} \cdot dx$  Using Simpson's

$\frac{1}{3}$  rule by taking  $h=1$ .

$\Rightarrow$  Here, Given  $f(x) = \frac{1}{1+x}$ ,  $a=0$ ,  
 $b=6$  and  $h=1$

$$\text{So, } n = \frac{b-a}{h} = \frac{6-0}{1} = 6$$

X	0	1	2	3	4	5	6
Y	1	0.5	0.333	0.25	0.2	0.167	0.143

By Simpson's  $\frac{1}{3}$  rule,

$$\int_0^6 \frac{1}{1+x} \cdot dx = \frac{h}{3} [ (Y_0 + Y_6) + 2(Y_2 + Y_4 + Y_6) + 4(Y_1 + Y_3 + Y_5) ]$$

$$= \frac{1}{3} [ (1 + 0.143) + 2(0.333 + 0.2 + 0.143) + 4(0.5 + 0.25 + 0.167) ]$$

$$\int_0^6 \frac{1}{1+x} \cdot dx = 1.959$$

2. Using Simpson's  $3/8$  rule evaluate  $\int_0^1 \frac{1}{1+x^2} dx$  by taking

$$h = 1/6$$

$\Rightarrow$  Here, Given  $f(x) = \frac{1}{1+x^2}$ ,  $a=0$ ,

$$b=1 \text{ and } h=1/6$$

$$\text{So, } n = \frac{b-a}{h} = \frac{1}{1/6} = 6$$

X	0	1/6	1/3	1/2	2/3	5/6	1
Y	1	0.9730	0.9	0.8	0.6923	0.5902	0.5

By Simpson's  $3/8$  rule,

$$\int_0^1 \frac{1}{1+x^2} dx = \frac{3h}{8} [Y_0 + Y_6 + 2(Y_3) + 3(Y_1 + Y_2 + Y_4 + Y_5)]$$

$$= \frac{3}{8 \times 6} [(1 + 0.5) + 2(0.8) + 3(0.6923 + 0.9 + 0.9730 + 0.5902)]$$

$$\int_0^1 \frac{1}{1+x^2} dx = 0.7854$$



3 Using Simpson's  $\frac{3}{8}$  rule evaluate  $\int_0^3 \frac{1}{1+x} \cdot dx$  by taking  $n=6$  and hence calculate  $\log 2$ .

$\Rightarrow$  Here, Given  $f(x) = \int_0^3 \frac{1}{1+x}$

$a = 0$ ,  $b = 3$  and  $n = 6$ .

$$\text{So, } h = \frac{b-a}{n} = \frac{3}{6} = 0.5$$

$x$	0	0.5	1	1.5	2	2.5	3
$y$	1	0.6666	0.5	0.4	0.333	0.2857	0.25

By Simpson's  $\frac{3}{8}$  rule,

$$\int_0^3 \frac{1}{1+x} \cdot dx = \frac{3(0.5)}{8} [(1+0.25) + 2(0.4) + 3(0.6666 + 0.5 + 0.3333 + 0.2857)]$$

$$\int_0^3 \frac{1}{1+x} \cdot dx = 1.3887$$

$$\begin{aligned} \Rightarrow \int_0^3 \frac{1}{1+x} \cdot dx &= \left| \log(1+x) \right|_0^3 \\ &= \log 4 \end{aligned}$$

4. Evaluate  $\int_{-2}^6 (1+x^2)^{3/2} dx$  by Simpson's  $1/3$  rule.

$\Rightarrow$  Here, Given,  $f(x) = (1+x^2)^{3/2}$ ,  
 $a = -2$ ,  $b = 6$  and  $n = 6$

$$\text{So, } h = \frac{b-a}{n} = \frac{6+2}{6} = 1.3333$$

x	-2	-2/3	2/3	2	10/3
y	11.1803	1.7360	1.7360	11.1803	42.1478

14/3	6
108.7088	225.0622

By Simpson's  $1/3$  rule,

$$\int_{-2}^6 (1+x^2)^{3/2} dx = \frac{h}{3} [(y_0 + y_6) + 2(y_2 + y_4) + 4(y_1 + y_3 + y_5)]$$

$$\int_{-2}^6 (1+x^2)^{3/2} = \frac{1}{8} (11.1803 + 225.0622) +$$

$$2(1.7360 + 42.1478) + 4(1.7360 +$$

$$11.1803 + 108.7088) ]$$

$$= \frac{1.333}{3} [236.2425 + 87.7676 +$$

$$480.5004]$$

$$\int_{-2}^6 (1+x^2)^{3/2} \cdot dx = 360.22$$

6 Evaluate  $\int_0^6 \frac{1}{1+x^2} \cdot dx$  using Weddle rule with  $n=6$ .

$\Rightarrow$  Here, Given  $f(x) = \frac{1}{1+x^2}$ ,  $a=0$ ,  
 $b=6$  and  $n=6$

$$h = \frac{b-a}{n} = \frac{6}{6} = 1$$

x	0	1	2	3	4	5	6
y	1	0.5	0.2	0.1	0.0588	0.0385	0.0270

By Weddle Rule,

$$\int_0^6 \frac{1}{1+x^2} \cdot dx = \frac{3h}{10} [y_0 + 5y_1 + y_2 + 6y_3 +$$

$$y_4 + 5y_5 + y_6]$$

$$= \frac{3}{10} (3.5 + 5(0.5))$$

$$\int_0^5 \frac{1}{1+x^2} \cdot dx = 1.3735$$

7 Evaluate  $\int_4^{5.2} \log x \cdot dx$  using  
Weddle rule with  $n=6$ .

$\Rightarrow$  Here, Given  $f(x) = \log x$ ,  $a=4$ ,  
 $b=5.2$  and  $n=6$ .

$$\text{So, } h = \frac{b-a}{n} = \frac{5.2-4}{6} = 0.2$$

x	4	4.2	4.4	4.6	4.8
y	1.3863	1.4351	1.4816	1.5260	1.5686

5	5.2
1.6094	1.6486

By using Weddle rule

$$\int_4^{5.2} \log x \cdot dx = \frac{3h}{10} [Y_0 + 5Y_1 + Y_2 + 6Y_3 + Y_4 + 5Y_5 + Y_6]$$

$$\int_4^{5.2} \log x \cdot dx = 1.8278$$

5 Evaluate  $\int_0^{\pi} \frac{\sin^2 x}{5+4\cos x} dx$  Using Simpson's  $3/8$  rule.

Here, Given  $f(x) = \frac{\sin^2 x}{5+4\cos x}$   
 $a = 0$ ,  $b = \pi$ ,  $h = \frac{\pi}{4}$ ,  $n = 4$

$x$	0	$\pi/4$	$\pi/2$	$3\pi/4$	$\pi$
$f(x)$	0	0.0639	0.2	0.2302	0

By Simpson's  $3/8$  rule

$$\int_0^{\pi} \frac{\sin^2 x}{5+4\cos x} \cdot dx = \frac{3h}{8} [y_0 + y_4 + 2y_2 + 3(y_1 + y_3)]$$

$$= \frac{3h}{8} [0 + 0.4604 + 0.9 + 7.917]$$

$$= 0.3688$$